


Saproxylic breeding sites for hoverflies (Diptera: Syrphidae): from artificial design to natural habitat management

Jeroen van Steenis



Journal van Syrphidae

An international journal on Syrphidae, “Journal van Syrphidae” (<http://zoobank.org/References/632F826B-7A33-4CA7-80D6-9B96135409C1>), founded in 2022 by the Syrphidae Foundation. *JvS* publishes original work in the form of articles, comments and symposium papers on all topics concerning Syrphidae (Insecta, Diptera) from all biogeographical regions.

Editor in Chief

Jeroen van Steenis. Syrphidae in trees, email: infojvs@syrphidaeintrees.com

Editorial Board

Leendert-Jan van der Ent, The Netherlands
Francis Gilbert, United Kingdom
Roger Morris, United Kingdom
Gerard Pennards, The Netherlands
Axel Ssymank, Germany
Tsung-Hsueh (Bill) Wu, Taiwan

Manuscripts

Manuscripts and other correspondences should be sent to the Editor in Chief.

The aim of the journal and “instructions for authors” can be found on the webpage of [JvS](#)

Publishing

Publication is through free online open access.

Indexing

The papers are supplied with DOI and Zoobank registration numbers. The authors are requested to register in ORCID or preferably in zoobank to get unique electronic identification numbers. See website for more information.

JvS is indexed in [Crossref](#).

Funding

JvS is financed through the [Syrphidae Foundation](#), and can only survive by donations of readers and other organisations or private persons gifts.

Organisations and private persons donating for the current Volume 2 are listed below.

veldshop.nl

(<https://www.veldshop.nl/en/>)



(<https://www.biomongol.org/>)

- J. Bisschop
- G.W.A. Pennards
- A.M. Ssymank
- J. van Steenis
- W. van Steenis
- M.P. van Zuijlen
- F.S. Zuidhoff

[[Donation button](#)]



Article

<https://doi.org/10.55710/1.DIOF2888>

<https://zoobank.org/References/1906E94E-6454-4E97-9A8C-197EB8CD5F05>

Saproxylic breeding sites for hoverflies (Diptera: Syrphidae): from artificial design to natural habitat management

Jeroen van Steenis

Syrphidae Foundation, Hof der Toekomst 48, 3823 HX Amersfoort, The Netherlands

e-mail: jvansteen@syphidaeintrees.com

<https://orcid.org/0000-0001-9231-1516>

Received: 28-11-2022

Accepted: 9-3-2023

Available online: 10-3-2023

Handling Editor: Axel Ssymank

Abstract. This paper introduces the importance of veteran trees, tree related microhabitats (TreMs) and their associated hoverfly (Diptera, Syrphidae) fauna. A broader perspective of creating larval habitat is discussed, based on published and novel insights. It focuses on hoverflies that specialise on veteran trees and reflects upon protection and management regimes to conserve veteran trees, TreMs and associated woody habitats. The lack of veteran trees breeding sites can be resolved by tree veteranisation or by using artificial breeding boxes. Whilst protection of veteran trees is essential, enhancement of open areas with flower resources is also vitally important for the survival of saproxylic hoverflies. The larval and adult ecology of only three out of the 134 known European saproxylic species are properly understood. Thus several suggestions are offered for future research aimed at a thorough understanding of the natural history of this unknown and ecologically relevant group of species. The list includes faunistic surveys and investigations into population dynamics, dispersal capacity and habitat preferences. Alongside this research there is a need to investigate the creation of breeding sites including veteranisation techniques and the use of breeding boxes.

Keywords. breeding boxes, hoverfly lagoons, protection management, Syrphidae in trees, TreMs, veteran trees

Citation: van Steenis J. 2023. Saproxylic breeding sites for hoverflies (Diptera: Syrphidae), from artificial design to natural habitat management. *Journaal van Syrphidae* 2(1): 1–22.

<https://doi.org/10.55710/1.DIOF2888>

Introduction

As adults, virtually all hoverfly species are known for their importance in pollinating crops and wildflowers all over the world (Ssymank *et al.* 2008; Inouye *et al.* 2015; Moquet *et al.* 2017; Doyle *et al.* 2020; Dunn *et al.* 2020) and the wider interest in this group of insects has

increased during the last decades (Fontaine *et al.* 2012; IUCN *et al.* 2022). On the contrary the larvae have a wide range of feeding strategies being from aquatic species to phytophagous, fungivores, and saproxylic species or others predacious on a wide range of preys (Rotheray & Gilbert 2011). Unfortunately, hoverflies are declining at an accelerating rate (Biesmeijer *et al.* 2006; van Eck 2016; Hallmann *et al.* 2017; Gatter *et al.* 2020; Hallmann *et al.* 2021; Morris & Ball 2021; Barendregt *et al.* 2022).

Based on the European Red List of hoverflies which was recently published the 134 species of saproxylic hoverflies represent 15% of the entire European hoverfly fauna of 892 species (Vujić *et al.* 2022). The number of saproxylic species in the different red-list categories, Data Deficient, Least Concern and Near Threatened are 9 (6,7%), 73 (54,5%) and 7 (5,2%) respectively. In the threatened categories Vulnerable, Endangered and Critically Endangered the number of species is 13 (9,7%), 30 (22,4%) and 2 (1,5%) respectively (Vujić *et al.* 2022a). The larvae of saproxylic Syrphidae occupy a wide variety of woody microhabitats in living trees, referred to as TreMs (tree related microhabitats, Fig. 1) and dead wood like stumps and logs (Rotheray 1993; Rotheray & Gilbert 2011).




Figure 1. Veteran tree with associated tree-related micro habitats (TreMs). Adjusted from Larrieu *et al.* 2017.

Adults of saproxylic species (Figs 2, 3) have been relatively well investigated, although faunistic information is still lacking for many species. The larvae, have had much less attention and worldwide only 8% of species had been described (Rotheray & Gilbert 2011), however a relevant work exist of several saproxylic species (e.g. Krivosheina & Mamayev 1962; Maibach & Goeldlin 1989; Rotheray 1991, 1993; Rotheray & Perry 1994; Rotheray & Stuke 1998; Rotheray & MacGowan 2000; Krivosheina 2001, 2004, 2018, 2019, 2020; Ricarte *et al.* 2007, 2009; van Steenis 2015; Orengo-Green *et al.* 2023).



Figure 2. Adult veteran tree specialized Syrphidae, with IUCN Red List category in parentheses. **A.** *Sphiximorpha subsessilis* (Illiger in Rossi, 1807) ♂ (LC), defending its territory on *Aesculus hippocastanum* with a sap run. Wijlre, The Netherlands. **B.** *Brachyopa testacea* (Fallén, 1817) ♂ (LC), defending its territory near pine logs (Fig. 6D). Teletskoye Lake, Artybash, Russia. **C.** *Milesia crabroniformis* (Fabricius, 1775) ♀ (LC), egg laying behaviour at a trunk base rot hole on *Platanus orientalis*. (Fig. 5D). Paphos forest, Appides stream, Cyprus. **D.** *Sphiximorpha petronillae* Rondani, 1850 ♀ (EN), egg laying behaviour on *Quercus pubescens* with sap runs (Fig. 5C). Kamenički Park, Novi Sad, Serbia.

Veteran trees are features of natural forests, but also occurring in parks or as single trees, containing many different types of TreMs (Fig. 1), such as trunk cavities, branch holes, exposed heartwood, dead branches and sap-runs (Read 2000, Larrieu *et al.* 2017). This variety of microhabitats has a significant impact on biodiversity, hosting about 2/3 of the terrestrial biodiversity, and enhancing ecosystem functioning (Speight 1989; Read 2000; Grove 2002; Ulyshen 2018; Wetherbee *et al.* 2021; Martin *et al.* 2022; Przepióra & Ciach 2022). Old growth forests provide significant carbon storage and have the capacity to withstand climate change better than young production forests (Hotchkiss 2020). These forests also have a positive effect on retaining water and thus regulation of rainfall and local reduction of temperature fluctuations (Rubino & McCarthy 2003; Sist *et al.* 2014; Hotchkiss 2020; Nolan

et al. 2020). Veteran trees also play a major role in pest management by providing habitat for pest predator species that reduce economic damage and thus decrease the demand for the use of harmful pesticides (Wetherbee *et al.* 2020).

The following exemplifies the importance of saproxylic insects “Saproxylic insects comprise the largest component of the biodiversity in terrestrial ecosystems. They are responsible for the mechanical breakdown of woody material both directly, by tunnelling and feeding in living trees that are decaying, snags (standing dead trees) and logs (fallen trees, portions of trunk and large branches), or indirectly, through symbiotic relationships with fungi and other microorganisms that humidify wood.” (Marcos-García *et al.* 2011). In natural and unmanaged forest TreMs reach a high diversity and are also more numerous compared to managed forests (Martin *et al.* 2022; Przepióra & Ciach 2022).




Figure 3. Adult veteran tree specialized Syrphidae, with IUCN Red List category in parentheses. **A.** *Callicera macquarti* Rondani, 1844 ♂ (EN), defending its territory on *Pinus brutia*. Paphos forest, Appides stream, Cyprus. **B.** *Brachypalpus valgus* (Panzer, 1798) ♂ (LC), defending its territory on *Salix alba* (Fig. 5B). University park. Novi Sad. Serbia. **C.** *Blera fallax* ♂ (LC), defending its territory on a pine stump (Fig. 6C). Teletskoye lake, Artybash, Russia. **D.** *Xylota pseudoignava* Mutin, 1984 ♂ (non European), defending its territory on a pine stump (Fig. 6C). Teletskoye lake, Artybash, Russia.

To establish conservation action plans, the biology of a species needs to be fully known, including its dispersal capacity, larval stages and the microhabitat preferences (Vujić *et al.* In press). So far, only three species, *Blera fallax* (Linnaeus, 1758), *Callicera rufa* Schummel, 1842 and *Hammerschmidtia ferruginea* (Fallén, 1817), have been included in species recovery plans (Rotheray & MacGowan 2000; Rotheray & Rotheray 2012; Rotheray 2013; Rotheray & MacGowan 2015). Many more such plans need to be included to ensure continuity and extent of the habitats of these species (e.g. Jukes 2009; IUCN *et al.* 2022).

This paper aims to give an overview of research to date on hoverflies that specialise in veteran trees. Other research on management to promote veteran trees and their associated TreMs will also be discussed, together with, proposals for future research.

References used in this paper are presented to stimulate further reading and only represent part of the exhaustive literature that can be found on this topic (van Steenis 2022).



Figure 4. Larval habitat veteran tree specialized Syrphidae. **A.** *Populus alba*, sap run. Fruška Gora, Stražilovo, Serbia. **B.** *Quercus infectoria*, trunk rot-hole. Tzelefos Bridge, Cyprus. **C.** Pine stump, sappy decay and trunk rot-hole. Teletskoye lake, Artybash, Russia. **D.** Pine logs. Teletskoye lake, Artybash, Russia.

Threatened veteran tree specialized Syrphidae found in the surroundings, IUCN category in parentheses.

A: *Brachyopa maculipennis* Thompson, 1980 (EN) and *B. plena* Collin, 1939 (NT).

B: *Ceriana glabosa* van Steenis & Ricarte in van Steenis *et al.* 2016 (EN).

C and D: *Blera eoa* (Stackelberg, 1928) (EN), *Callicera aenea* (Fabricius, 1777) (VU), *Chalcosyrphus femoratus* (Linnaeus, 1758) (VU), *C. rufipes* (Loew, 1873) (EN), *Sphecomyia vespiformis* (Gorski, 1852) (NT), *Temnostoma angustistriatum* Krivosheina, 2002 (EN) and *T. apiforme* (Fabricius, 1794) (NT).

Creating breeding sites for Syrphidae

Creation of hoverfly breeding sites has happened since humans started to shape the environment to meet its own needs. Examples include felling wood for cooking, creating tools and housing and thus creating coppice and also stumps and logs. These actions were not directly intended to create valuable habitat but have influenced the occurrence of saproxylic

species (Rösch 2012; Fazan *et al.* 2020). More recently, creation of artificial breeding places for birds (Onur Erman 2014) and bats (Rueegger 2016) has become standard conservation tools, whilst bee-hotels have become a popular addition to the list of artificial breeding sites (MacIvor 2017). Many papers have been published about the differences between natural tree-holes, man-made tree-holes and artificial breeding boxes in a wide variety of research setups. These investigations all demonstrate the value of breeding boxes and the composition of the breeding medium that is used (Jansson *et al.* 2009; Carlsson *et al.* 2016; Griffiths *et al.* 2018; Dulisz *et al.* 2022). Creation of artificial breeding sites for Syrphidae was first investigated by Maibach & Goedlin (1989) and MacGowan (1994), followed by many attempts to enhance the effectiveness of the artificial breeding medium (e.g. MacGowan & Rotheray 2007; Rotheray & MacGowan 2015; Foster & Leach 2021; RZSS 2022). Several of these attempts only had very limited results, mostly attracting *Myathropa florea* (Linnaeus, 1758) or an inability to maintain a population with larvae dying before reaching maturity (Rotheray 2004; Schmid & Moertelmaier 2007; van Steenis 2015; Wetherbee *et al.* 2022). In other cases it was possible to maintain moisture over several years and produce continuing generations of saproxylic Syrphidae (Rotheray 2013; Taylor *et al.* 2021). These examples suggest that larval biology of saproxylic species requires more detailed understanding microhabitat preferences and the specific nutritional, physical and chemical composition of its breeding medium, thus, much more research is needed (Sánchez-Galván *et al.* 2014; Ulyshen 2018; IUCN *et al.* 2022; Wetherbee *et al.* 2022).

Artificially created breeding sites need maintenance (Lindemayer *et al.* 2009) with special attention paid to ensure suitable humidity (Rotheray 2004). Personal experience suggests that this process is demanding. Artificial sites are therefore imperfect for species protection (Jansson *et al.* 2009). Instead, they should be seen as a means of enhancing short term survival, thus making research possible without risking local extinction of the species under investigation. A more sustainable way of protecting saproxylic Syrphidae species depends upon habitat protection of larval and adult and through preservation of veteran trees, their associated TreMs and food sources for the adults.

Practices to sustain saproxylic species

Forestry management

In forestry practice clear-felling large areas is commonplace, but it often leads to the creation of stumps, snags and living tall stumps that maintain, at least some of, the habitat utilised by saproxylic faunas (e.g. Jonsell *et al.* 2004; Sandström *et al.* 2019). Most of the research conducted so far has involved Coleoptera. It shows that the most important factor in maintaining high biodiversity of saproxylics is the creation of tall and low stumps (e.g. Jonsell & Weslien 2003; Lindhe & Lindelöw 2003). Felling, preferably during winter, or at least well before the start of the season has a positive impact on beetle abundance (Foit 2012). Species abundance and composition does not differ much between low or tall pine stumps (Hjälten *et al.* 2010); however, in broadleaved trees, it seems that tall stumps represent a more favourable habitat (Jonsell *et al.* 2004; Lindhe & Lindelöw 2004). Leaving long logs of either tree species increases biodiversity even more (Jonsell & Weslien 2003). Saproxylic beetles and Syrphidae respond differently to forestry management. For example, longhorn beetles prefer dense oak-dominated areas with a high volume of snags and veteran trees, while saproxylic hoverflies are more limited to open-stands with large veteran trees and a well-developed, species rich herb layer (Fayt *et al.* 2006; Nol *et al.* 2006). Diptera and especially Syrphidae larvae need moisture to develop like dendrotlems (water filled rot-holes), here differentiated

from dry rot-holes (Fig. 1), sap-runs and moist decaying stumps and logs.




Figure 5. Larval habitat veteran tree specialized Syrphidae. **A.** *Salix alba*, limb breakage and sap run. University park, Novi Sad, Serbia. **B.** *Salix alba*, sap runs, cracks and trunk base rot-holes. University park, Novi Sad, Serbia. **C.** *Quercus pubescens*, several limb breakages, sap runs and trunk rot-holes. Kamenički Park, Novi Sad, Serbia. **D.** *Platanus orientalis*, trunk base rot-hole. Paphos forest, Appides stream, Cyprus.

Threatened veteran tree specialized Syrphidae found in the surroundings, IUCN category in parentheses.

A and B: *Brachyopa maculipennis* (EN).

C: *Brachyopa grunewaldensis* Kassebeer, 2000 (EN), *B. silviae* Doczkal & Dziock, 2004 (EN), *Brachypalpus* aff *valgus* (possibly CR) and *Sphiximorpha petronillae* (EN).

D: *Callicera macquarti* (EN).

Sap-runs and partially water submerged logs are almost exclusively inhabited by larvae of this order (Maibach & Goeldlin de Tiefenau 1992; Jukes 2009; van Eck 2016; van Eck *et al.* 2016;

Wolton & Luff 2016). Thus, not only veteran trees but the entire forest habitat needs conservation (Hotchkiss 2020). Several of the important tree species such as oak, beech and pine can take more than 100 years to become a multiple TreMs tree (Hövmeyer & Schauer mann 2003; Drobyshchev *et al.* 2008; Piovesan & Biondi 2021) able to host a wide variety of saproxylic insects, while they can reach an age of more than 1000 years (Gandy 2019; van Steenis *et al.* 2019; Nolan *et al.* 2020). Also, the time span in which wet decay, a suitable medium for the larvae of many saproxylic Syrphidae, will be formed or will last depends also on the actual tree species (Rotheray & Gilbert 2011). Conversely, fast growing trees such as birch (*Betula* spp.), poplars and aspen (*Populus* sp.) mature at a faster rate (Nieuwenhuis & Barrett 2002; Ninufu 2008; Myking *et al.* 2011; Caudullo & de Rigo 2016) and logs can be productive as breeding sites within just 20–50 years when diameters have reached 10–25 cm (Rotheray 1993; Krivosheina 2004; Drees 1999; Rotheray *et al.* 2008). There is a succession in the species composition in decaying logs in which the first years sappy decay develops under intact bark, which turns into bark losing hardwood logs and eventually become wet soft wood decay. The sappy decay may persist from 2–7 years after which the hardwood period will last an additional 2–5 years (Derksen 1941; Hövmeyer & Schauer mann 2003; Rotheray *et al.* 2008; Krivosheina 2018, 2019).



Figure 6. Adult habitat veteran tree specialized Syrphidae. **A.** Kamenički Park, Novi Sad, Serbia. **B.** Flowering *Hedera helix*. Imbros Gorge, Crete, Greece. **C.** Flower rich forest glade. Riikanmaa, Joutseno, Finland. **D.** Flower rich forest glade. Strandzha; Sinemorets, Bulgaria. Threatened veteran tree specialized Syrphidae found in the surroundings, IUCN category in parentheses. A: see Fig. 5C.

B: *Callicera macquarti* (EN), *C. rufa* (VU), *C. spinolae* Rondani, 1844 (VU) and *Milesia cretica* Bot & van Steenis in Bot *et al.* 2022 (possibly VU).

C: *Temnostoma apiforme* (NT) and *Temnostoma sericomylaeforme* (Porschinsky, 1887) (VU).

D: *Chalcosyrphus pannonicus* (Oldenberg, 1916) (EN) and *Psilota nana* Smit & Vujić, 2008 (EN).

Veteran trees and their remnant stumps and logs play a key role in the survival of saproxylic species (Figs 1, 4, 5), however, in Syrphidae much more is needed than the protection of these trees. Without adequate flowering resources (Fig. 6) the adults will not be able to mature, nor to develop their eggs for which nectar and pollen are essential (Szymank & Gilbert 1993; Moquet *et al.* 2017; Nicholas *et al.* 2018). The problem with forestry conservation is that a closed canopy diminishes the development of TreMs on veteran trees (Miklín *et al.* 2018) and also hinders the growth of flower resources for the adult Syrphidae (van Steenis 2016; Hotchkiss 2022).

Preservation of veteran trees outside forests

Saproxylic Syrphidae are mostly associated with forest habitats, but urban parks and even single (hedgerow) trees can harbour many saproxylic Syrphidae (Coe 1953; Ahnlund 1996; Andersson 1999; Sjuts 2004; van Steenis *et al.* 2019). As such, they have the potential to form important islands and steppingstones in an otherwise forest free environment. The protection of forest habitats, trees in city parks and single trees is therefore of considerable importance. Trees in floodplain forests (Figs 5A, 5B), in city parks (Fig. 5C) or single trees also known as trees outside of forests (TOF) hold many TreMs with a large diversity of saproxylic species and need protection (Coe 1953; Andersson 1999; Schlaghamerský 2003; de Foresta 2013; Alexander *et al.* 2015; Miklín *et al.* 2017; Gandy 2019; van Steenis *et al.* 2019). Protection management should thus be directed to ensuring the continued presence of veteran trees, open areas and enhancing hydrology (Peterken 2002; Davies *et al.* 2008; Hotchkiss 2022).

Breeding boxes

In areas where there are very few old or veteran trees, the use of breeding boxes could help to overcome the lack of suitable breeding habitat. It should be noted that the maintenance of these breeding boxes is time consuming and it should only be used as short-term solution or in enhancing populations for research purposes. The effectiveness of these artificial breeding sites has never been fully tested, although several efforts have been made to produce these so-called breeding boxes with varying degrees of success (Maibach & Golecllin 1992; Rotheray 2004; Schmid & Moertelmaier 2007; Rotheray & MacGowan 2015).

Ideally, most effort should focus upon veteranisation of trees instead of maintaining less reliable and more time consuming artificial habitats. Nevertheless, creating easily made and maintained hoverfly lagoons for saprophagous aquatic Syrphidae has been very successful in Great Britain (Buzz Club 2018). The breeding medium used consisted of different decaying matter, mostly grass (Fig. 7A), but also decaying vegetables such as potatoes and carrots, fresh sawdust, or droppings of different small or large herbivores. In my own experience, most species were attracted to hoverfly lagoons positioned on the ground and filled with decaying vegetables. The lagoons and pet-bottles with sawdust attached to trees at different heights only attracted *Myathropa florea*. When a more robust breeding box (Figs 7B, 7C) was trialled, incorporating different sawdust mixtures, again *M. florea* was almost exclusively the occupant. Breeding medium mixture will probably make a great difference in attracting different species and this needs to be investigated in greater detail.



Figure 7. Artificial breeding sites. **A.** Hoverfly lagoon for aquatic Syrphidae (from the Buzz Club, 2018). **B, C.** Breeding boxes from “Syrphidae in trees” for saproxylic Syrphidae. **B.** Breeding box in pine, Soest, The Netherlands. **C.** From left to right: “internal” Pet-bottle surrounded by saw-dust (back-panel not mounted); saw-dust filled box (back panel not mounted); lateral view; frontal view, with closed back panel, before hanging the box in the tree a second Pet-bottle filled with saw-dust and water was placed into the “internal” Pet-bottle.

Recommendations for preservation of saproxylic species

In the literature, different recommendations are suggested for conservation of saproxylic species. In aiming for 4–16 veteran trees per hectare there is a need to ensure a supply 5–20 young trees per hectare: ‘legacy trees’ that will become veteran trees. They should be protected from felling or other damage causing decline in the number of TreMs and these trees should be marked in an unambiguous way, not like in Fig. 8. In some cases, intentionally damaging trees might enhance the formation of TreMs (e.g. Lewis 1998; Jonsell *et al.* 2004). Maintaining or even creating “open” space such as forest edges, glades or places around legacy trees is also needed to increase the species richness of adult feeding sources like flowering herbs, shrubs and trees as in Syrphidae adult flower visiting occurs close to the larval tree-related habitats, typically on the forest margin itself and up to 100–200 m into open meadows (Ssymank 1991, 2001; Falk 2021). About 10–15% of trees should be decaying snags or stumps and logs. Other action might include blocking drainage ditches and leaving

trees and logs in certain parts of the river system (Fig. 8C), in which this last action has a positive effect on *Chalcosyrphus eunotus* (Loew, 1873) (Read 2000; Boesch *et al.* 2007; Jukes 2009; Soszyńska-Maj *et al.* 2009; Sebek *et al.* 2013; Hotchkiss 2022). However, these protection measurements are general, and little is known about the true habitat preferences of many saproxylic Syrphidae and for this further research is needed.



Figure 8. Larval habitat veteran tree specialized Syrphidae. **A.** *Pinus sylvestris*, dendrotelm fork split. Lange Duinen, Soest, The Netherlands. **B.** Same tree cut down for “nature conservation” despite being marked with red and white tape. Both the stump and log were later removed and destroyed. **C.** Rivulet with dead trees and partly water submerged logs. “Bunderbos, Zuid-Limburg, The Netherlands.

Threatened veteran tree specialized Syrphidae found in the surroundings, IUCN category in parentheses.

A, B. *Callicera rufa* (VU).

C. *Chalcosyrphus eunotus* (VU).

Discussion

Remarks on future research

The following suggestions offer ideas on future research needed to investigate the habitat preferences of saproxylic Syrphidae in order to practical habitat management for saproxylic hoverflies. Some research has already been done on these issues and examples of references

are given after each research task. Extensive research has been done concerning Coleoptera as regards to the micro-habitat requirements, dispersal capacity and the effects of creating breeding sites and some of these references will sometimes be given here alongside those dealing with saproxylic Syrphidae.

1. Review the knowledge on faunistics of the target species or genera (Rotheray, 2013; Pérez-Bañón *et al.* 2016; van Steenis *et al.* 2020; Falk 2021).
2. Search for additional populations of the species through targeted search for suitable habitat, host trees or accompanying species such as the European velvety tree ant (*Liometopum microcephalum* (Panzer, 1798)) in the case of *Sphiximorpha petronillae* (Birtele 2003; Del Toro *et al.* 2009; Sebel *et al.* 2013; Mei 2016; Miklín *et al.* 2017; Griffiths *et al.* 2018; Tăușan 2018; van Steenis *et al.* 2019; Bütler *et al.* 2020; Petermann & Gossner 2022).
3. Investigate dispersal capacity as vital factor for implementing habitat restoration efforts (Reenema 2000; Rotheray *et al.* 2014; Komonen & Müller 2018, Martínez-Pérez *et al.* 2022).
4. Investigate the influence of flower resources on fitness and fecundity of adults and a possible link to larval survival rates (Ssymank 1991; Branquart & Hemptinne 2000; van Rijn & Wäckers 2016).
5. Undertake practical investigation into veterinisation of trees by creating dendrotelm trunk holes, “open” dendrotelm (base or fork-split) holes, sap-runs, broken off branches with wound healing, high and low stumps, snags, logs investigating the effects of size, humidity etc. (MacGowan 1994; Cavalli & Mason 2003; Rotheray & Rotheray 2012).
6. Study the effect of grazing on veterinization or destruction of the TreMs (Rotheray & MacGowan 2000; Stiven 2009; Pérez-Bañón *et al.* 2016; Ramón Arévalo *et al.* 2021).
7. Review literature or targeted field research on xylobiont insects creating or enhancing breeding habitat for Syrphidae (Quinto *et al.* 2012; Sánchez-Galván *et al.* 2014; Krivosheina 2019, 2020).
8. Creating breeding boxes to be used during population dynamic studies or in young growth forests to establish jumping populations to and from isolated wild populations (Maibach & Goeldlin 1992; Rotheray 2004; Carlsson *et al.* 2006; Schmid & Moertelmaier 2007; Jansson *et al.* 2009; van Steenis 2015; Landvik *et al.* 2016).
9. Investigate how to incorporate nature conservation actions with raising awareness to the people through social media and citizen science (Read 2000; Flint *et al.* 2009; Thorn *et al.* 2020; Falk 2021).

Acknowledgments

Ante Vujić (Novi Sad, Serbia) is greatly acknowledged for his work on protecting Syrphidae and initiating many projects, not least the Conservation Action Plan for hoverfly species that specialise in veteran trees in Europe. Marija Miličić (Novi Sad, Serbia) gave valuable comments on the draft version of this paper, Axel Ssymank (Wachtberg, Germany) provided additional literature and Roger Morris (Mitcham, UK) checked the correct use of the English language.

References

- Alexander K.N.A., Bengtsson V.J., Jansson N. & Smith J.P. 2016. The role of trees outside woodlands in providing habitat and ecological networks for saproxylic invertebrates Part 1 Designing a field study to test initial hypotheses. Natural England, Worcester. 108 pp. <http://publications.naturalengland.org.uk/publication/4828234842112000>
- Andersson H. 1999. Rödlistade eller sällsynta evertebrater knutna till ihåliga, murkna eller savande träd samt träd svampar i Lunds stad. [Red-listed or rare invertebrates associated with hollow, rotting, or sapping trees or polypores in the town of Lund]. Entomologisk Tidskrift 120 (4): 169–183. https://www.sef.nu/download/entomologisk_tidskrift/et_1999/1999%20169_183.pdf
- Ahnlund H. 1996. Vedinsekter på en sörmländsk aspstubbe [Saproxylic insects on a Swedish dead aspen]. Entomologisk Tidskrift. 117 (4): 137–144. https://www.sef.nu/download/entomologisk_tidskrift/et_1996/ET1996%20137-144w.pdf
- Barendregt A., Zeegers T., van Steenis W. & Jongejans E. 2022. Forest hoverfly community collapse: Abundance and species richness drop over four decades. Insect Conservation & Diversity 15(5): 510–521. <https://doi.org/10.1111/icad.12577>
- Biesmeijer J.C., Roberts S.P.M., Reemer M., Ohlemüller R., Edwards M., Peeters T., Schaffers P., Potts S.G., Kleukers R., Thomas C.D., Settele J. & Kunin W.E. 2006. Parallel Declines in Pollinators and Insect-Pollinated Plants in Britain and the Netherlands. Science 313 (5785): 351–354. <https://doi.org/10.1126/science.1127863>
- Birtele D. 2003. The succession of saproxylic insects in dead wood: a new research method. In: Mason F., Nardi G. & Tisato M. (eds) Dead wood: a key to biodiversity. LIFE nature project NAT/IT/99/6245. Montova, Italy, pp 91–93.
- Boesch A., Pellet J. & Maibach A. 2007. Reconversion de populicatures et biodiversité. [Conversion of poplar plantations and biodiversity]. Schweizerische Zeitschrift für Forstwesen 158 (10): 323–330. <https://doi.org/10.3188/szf.2007.0323>
- Branquart E. & Hemptinne J-L. 2000. Development of ovaries, allometry of reproductive traits and fecundity of *Episyrphus balteatus* (Diptera: Syrphidae). European Journal of Entomology 97 (2): 165–170. <https://doi.org/10.14411/eje.2000.031>
- Bütler R., Lachat T., Krumm F., Kraus D. & Larrieu L. 2020. Field Guide to Tree-related Microhabitats. Descriptions and size limits for their inventory. Birmensdorf, Swiss Federal Institute for Forest, Snow and Landscape Research WSL. 59 pp. https://www.dora.lib4ri.ch/wsl/islandora/object/wsl%3A25942/datastream/PDF/B%C3%BCtler-2020-Field_guide_to_tree-related_microhabitats.-%28published_version%29.pdf
- Buzz Club 2018. The science behind wildlife gardening. Hoverfly Lagoons. School of Life Sciences, University of Sussex, Falmer, Brighton. <https://www.thebuzzclub.uk/hoverfly-lagoons> [accessed on 16-11-2022]
- Carlsson S., Bergman K-O., Jansson N., Ranius T. & Milberg P. 2016. Boxing for biodiversity: evaluation of an artificially created decaying wood habitat. Biodiversity and Conservation 25: 393–405. <https://doi.org/10.1007/s10531-016-1057-2>
- Caudullo G. & de Rigo D. 2016. *Populus tremula* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayaz J., de Rigo D., Caudullo G., Houston Durrant T. & Mauri A. (eds), European Atlas of Forest Tree Species. Publ. Off. EU, Luxembourg: 138–139. https://forest.jrc.ec.europa.eu/media/atlas/Populus_tremula.pdf
- Cavalli R. & Mason F. 2003. Techniques for reestablishment of dead wood for saproxylic fauna conservation. LIFE nature project NAT/IT/99/6245 Bosco della Fontana (Mantova, Italy). Gianluigi Arcari. 112 pp.

- Coe R.L. 1953. *Mallota cimbiciformis* Fallén (Diptera: Syrphidae) breeding in Hyd Park, London. Its larva and puparium compared with those of *Eristalis tenax* L. *Myiathropa florea* L. and *Helophilus* spp. Entomologist's Gazette 4: 282–286.
- Davies Z.G., Tyler C., Stewart G.B. & Pullin A.S. 2008. Are current management recommendations for saproxylic invertebrates effective? A systematic review. Biodiversity and Conservation 17(1): 209–234. <https://doi.org/10.1007/s10531-007-9242-y>
- de Foresta H., Somarriba E., Temu A., Boulanger D., Feuilly H. & Gauthier M. 2013. Towards the assessment of trees outside forests. Resources Assessment Working Paper 183. FAO Rome. 335 pp. <https://www.fao.org/3/aq071e/aq071e.pdf>
- Del Toro I., Pacheco J.A. & Mackay W.P. 2009. Revision of the Ant Genus *Liometopum* (Hymenoptera: Formicidae). Sociobiology. 53(2A): 296–369.
- Derksen W. 1941. Die Succession der Pterygoten Insekten im Abgestorbenen Buchenholz. [The succession of pterygote insects in dead beech wood]. Zeitschrift für Morphologie und Ökologie der Tiere 37: 683–734. <https://doi.org/10.1007/BF00437762>
- Drees M. 1999. Erfahrungen mit der Aufzucht von *Temnostoma bombylans* (Fabricius, 1805) und *T. vespiforme* (Linnaeus, 1758) aus den Larven (Diptera, Syrphidae). [Rearing observations on the larvae of *Temnostoma bombylans* (Fabricius, 1805) and *T. vespiforme* (Linnaeus, 1758) (Diptera, Syrphidae)]. Volucella 4 (1/2): 121–126. https://www.zobodat.at/pdf/Volucella_4_0121-0126.pdf
- Drobyshev I., Niklasson M., Linderson H., Sonesson K., Karlsson M., Nilsson S.G. & Lanner J. 2008. Lifespan and mortality of old oaks - Combining empirical and modelling approaches to support their management in Southern Sweden. Annals of Forest Science 65(4): 1–12. <https://doi.org/10.1051/forest:2008012>
- Doyle T., Hawkes W.L., Massy R., Powney G.D., Menz M.H. & Wotton K.R. 2020. Pollination by hoverflies in the Anthropocene. Proceedings of the Royal Society B 287(1927): 20200508. <https://doi.org/10.1098/rspb.2020.0508>
- Dulisz B., Stawicka A.M., Knozowski P., Diserens T.A. & Nowakowski J.J. 2022. Effectiveness of using nest boxes as a form of bird protection after building modernization. Biodiversity and Conservation 31: 277–294. <https://doi.org/10.1007/s10531-021-02334-0>
- Dunn L., Lequerica M., Reid C.R. & Lattya T. 2020. Dual ecosystem services of syrphid flies (Diptera: Syrphidae): pollinators and biological control agents. Pest Management Science 76(6): 1973–1979. <https://doi.org/10.1002/ps.5807>
- Falk S. 2021. A review of the pollinators associated with decaying wood, old trees and tree wounds in Great Britain. Woodland Trust, Grantham, Lincolnshire, Great Britain. 143 pp. <http://doi.org/10.13140/RG.2.2.31078.14408>
- Fayt P., Dufrêne M., Branquart E., Hastir P., Pontégnie C., Henin J-M. & Versteirt V. 2006. Contrasting responses of saproxylic insects to focal habitat resources: the example of longhorn beetles and hoverflies in Belgian deciduous forests. Journal of Insect Conservation 10: 129–150. <https://doi.org/10.1007/s10841-006-6289-0>
- Fazan L., Song Y-G. & Kozłowski G. 2020. The Woody Planet: From Past Triumph to Manmade Decline. Plants 9(11): 1593. <https://doi.org/10.3390/plants9111593>
- Flint C.B., McFarlane B. & Müller M. 2009. Human dimensions of forest disturbance by insects: An international synthesis. Environmental Management 43: 1174–1186. <https://doi.org/10.1007/s00267-008-9193-4>
- Foit J. 2012. Early-arriving saproxylic beetles developing in Scots pine stumps: effects of felling type and date. Journal of Forest science 58(11): 503–512. <https://doi.org/10.17221/46/2012-JFS>

- Fontaine B., van Achterberg K., Alonso-Zarazaga M.A., Araujo R., Asche M., Aspo H., Aspo U., Audisio P., Aukema B., Bailly N., Balsamo M., Bank R.A., Belfiore C., Bogdanowicz W., Boxshall G., Burckhardt D., Chylarecki P., Deharveng L., Dubois A., Enghoff H., Fochetti R., Fontaine C., Gargominy O., Soledad Gomez Lopez M., Goujet D., Harvey M.S., Heller K-G., van Helsdingen P., Hoch H., de Jong Y., Karsholt O., Los W., Magowski W., Massard J.A., McInnes S.J., Mendez L.F., Mey E., Michelsen V., Minelli A., Nieto Nafria J.M., van Nieukerken E.J., Pape T., de Prins W., Ramos M., Ricci C., Roselaar C., Rota E., Segers H., Timm T., van Tol J. & Bouchet P. 2012. New Species in the Old World: Europe as a Frontier in Biodiversity Exploration, a Test Bed for 21st Century Taxonomy. PLoS ONE 7(5): e36881. <https://doi.org/10.1371/journal.pone.0036881>
- Foster R. & Leach J. 2020. Furry Pine Hoverfly discoveries in the North Midlands. Hoverfly Newsletter 68: 3–4. <https://dipterists.org.uk/sites/default/files/pdf/Hoverfly%20Newsletter%2068.pdf>
- Foster R. & Leach J. 2021. *Chalcosyrphus nemorum* larvae in a beech stump hoverfly lagoon. Hoverfly Newsletter 69: 5–9. <https://dipterists.org.uk/sites/default/files/pdf/Hoverfly%20Newsletter%2069%20abridged%20version.pdf>
- Gandy M. 2019. The fly that tried to save the world: Saproxylic geographies and other-than-human ecologies. Transactions of the Institute of British Geographers 44(2): 392–406. <https://doi.org/10.1111/tran.12281>
- Gatter W., Ebenhöf H., Kima R., Gatter W. & Scherer F. 2020. 50-jährige Untersuchungen an migrierenden Schwebfliegen, Waffenfliegen und Schlupfwespen belegen extreme Rückgänge (Diptera: Syrphidae, Stratiomyidae; Hymenoptera: Ichneumonidae). [50-year studies of migrating hoverflies, soldier flies and parasitic wasps show extreme declines (Diptera: Syrphidae, Stratiomyidae; Hymenoptera: Ichneumonidae)]. Entomologische Zeitschrift 130 (3): 131–142. https://randecker-maar.de/wp-content/uploads/2020/10/2020_50-j%C3%A4hr.-Untersuchungen-an-Schwebfliegen-belegen-extreme-R%C3%BCckg%C3%A4nge.pdf
- Griffiths S.R., Lentini P.E., Semmens K., Watson S.J., Lumsden L.F. & Robert K.A. 2018. Chainsaw-Carved Cavities Better Mimic the Thermal Properties of Natural Tree Hollows than Nest Boxes and Log Hollows. Forests 9(5): 235. <https://doi.org/10.3390/f9050235>
- Hallmann C.A., Sorg M., Jongejans E., Siepel H., Hofland N., Schwan H., Stenmans W., Müller A., Sumser H., Hörren T., Goulson D. & de Kroon H. 2017. More than 75 percent decline over 27 years in total flying insect biomass in protected areas. PLoS ONE 12(10): e0185809. <https://doi.org/10.1371/journal.pone.0185809>
- Hallmann C.A., Ssymank A., Sorg M., de Kroon H. & Jongejans E. 2021. Insect biomass decline scaled to species diversity: General patterns derived from a hoverfly community. Proceedings of the National Academy of Sciences 118(2): e2002554117 <https://doi.org/10.1073/pnas.2002554117>
- Hartley J.C. 1961. A taxonomic account of the larvae of some British Syrphidae. Proceedings of the zoological Society London 136(4): 505–573. <https://doi.org/10.1111/j.1469-7998.1961.tb05891.x>
- Hotchkiss A. 2020. Ancient Woodland Restoration – Phase three: maximising ecological integrity. Woodland Trust Practical Guidance. The Woodland Trust, Grantham, UK. 48 pp. <https://www.woodlandtrust.org.uk/media/49219/ancient-woodland-restoration-maximising-ecological-integrity.pdf>
- Hövmeyer K. & Schauerermann J. 2003. Succession of Diptera on dead beech wood: A 10-year study. Pedobiologia 47: 61–75. <https://diptera.info/downloads/Hovmeyer.pdf>

- Inouye D.W., Larson B.M.H., Ssymank A. & Kevan P.G. 2015. Flies and Flowers III: Ecology of foraging and pollination. *Journal of Pollination Ecology* 16(16): 115–133. [https://doi.org/10.26786/1920-7603\(2015\)15](https://doi.org/10.26786/1920-7603(2015)15)
- IUCN SSC & HSG/CPSG. 2022. European Hoverflies: Moving from Assessment to Conservation Planning. Conservation Planning Specialist Group, Apple Valley, MN, USA. 84 pp. <https://doi.org/10.2779/359875>
- Jansson N., Ranius T., Larsson A. & Milberg P. 2009. Boxes mimicking tree hollows can help conservation of saproxylic beetles. *Biodiversity and Conservation* 18: 3891–3908. <https://doi.org/10.1007/s10531-009-9687-2>
- Jonsell M. & Weslien J. 2003. Felled or standing retained wood – it makes a difference for saproxylic beetles. *Forest Ecology and Management* 175(1–3): 425–435. [https://doi.org/10.1016/S0378-1127\(02\)00143-3](https://doi.org/10.1016/S0378-1127(02)00143-3)
- Jonsell M., Nittérus K. & Stighäll K. 2004. Saproxylic beetles in natural and man-made deciduous high stumps retained for conservation. *Biological Conservation* 118(2): 163–173. <https://doi.org/10.1016/j.biocon.2003.08.017>
- Jukes A. 2009. *Chalcosyrphus eunotus* a red data book hoverfly. Its status, distribution, ecology and conservation. Staffordshire Wildlife Trust (SWT). 21 pp. <https://www.staffs-wildlife.org.uk/sites/default/files/2018-12/Chalcosyrphus%20eunotus.pdf>
- Komonen A. & Müller J. 2018. Dispersal ecology of deadwood organisms and connectivity conservation. *Conservation Biology* 32(3): 535–545. <https://doi.org/10.1111/cobi.13087>
- Krivosheina M.G. 2001. Notes on the biology of palaeartic flies of the genera *Chalcosyrphus* Curran and *Xylota* Meigen (Diptera, Syrphidae), with the description of immature stages of *Xylota atricoloris* Mutin, 1987. *International Journal of Dipterological Research* 12(3): 165–172.
- Krivosheina M.G. 2004. A Contribution to the Biology of Flower Flies of the Genus *Temnostoma* (Diptera, Syrphidae) with Description of Larvae of Four Species. *Zoologicheskii Zhurnal* 82(1): 44–51.
- Krivosheina N.P. 2018. Habitat associations of the Larvae of the genus *Sphegina* Meigen, 1822 (Diptera, Syrphidae) with Xylobiont Insects. *Entomological Review* 98(6): 709–713. <https://doi.org/10.1134/S0013873818060076>
- Krivosheina N.P. 2019. Biotopic relations of flower-fly larvae of the genus *Brachyopa* Meigen, 1822 (Diptera, Syrphidae) and other Xylobiont insects. *Zoologicheskii Zhurnal* 98(9): 1063–1071 [In Russian with English summary]
- Krivosheina N.P. 2020. Biotopic associations of the larvae of the Hoverfly tribe Xylotini (Diptera, Syrphidae: Eristalinae) with Xylobiont Insects. *Entomological Review* 100 (2): 200–212. <https://doi.org/10.1134/S0013873820020086>
- Krivosheina N.P. & Mamayev B.M. 1962. Larvae of the European species of the genus *Temnostoma* (Diptera, Syrphidae). *Entomological Review* 41: 570–575.
- Landvik M., Niemelä P. & Roslin T. 2016. Mother knows the best mould: an essential role for non-wood dietary components in the life cycle of a saproxylic scarab beetle. *Oecologia* 182: 163–175. <https://doi.org/10.1007/s00442-016-3661-y>
- Larrieu L., Paillet Y., Winter S., Bütler R., Kraus D., Krumm F., Lachat T., Michel A.K., Regnery B. & Vandekerckhove K. 2018. Tree related microhabitats in temperate and Mediterranean European forests: A hierarchical typology for inventory standardization. *Ecological Indicators* 84:194–207. <https://doi.org/10.1016/j.ecolind.2017.08.051>
- Lewis J. 1998. Creating Snags and Wildlife Trees in Commercial Forest Landscapes. *Western Journal of Applied Forestry* 13(3): 97–101. <https://doi.org/10.1093/wjaf/13.3.97>
- Lindhe A. & Lindelöw Å. 2003. Cut high stumps of spruce, birch, aspen and oak as breeding substrates for saproxylic beetles. *Forest Ecology and Management* 203(1–3): 1–20. <https://doi.org/10.1016/j.foreco.2004.07.047>

- Lindenmayer B.B., Welsh A., Donnelly C., Crane M., Michael D., Macgregor C., McBurney L., Montague-Drake R. & Gibbons P. 2009. Are nest boxes a viable alternative source of cavities for hollow-dependent animals? Long-term monitoring of nest box occupancy, pest use and attrition. *Biological Conservation* 142 (1): 33–42. <https://doi.org/10.1016/j.biocon.2008.09.026>
- MacGowan I. 1994. Creating breeding sites for *Callicera rufa* Schummel (Diptera, Syrphidae) and a further host tree. *Dipterists Digest* 1: 6–8.
- MacGowan I. & Rotheray G.E. 2007. *Callicera rufa* Schummel Diptera, Syrphidae status and trends an update. *Dipterists Digest Second Series* 132: 113–118
- MacIvor J.S. 2017. Cavity-nest boxes for solitary bees: a century of design and research. *Apidologie* 48: 311–327. <https://doi.org/10.1007/s13592-016-0477-z>
- Maibach A.M. & Goeldlin de Tiefenau P. 1989. *Mallota cimbiciformis* (Fallén) nouvelle pour la faune de Suisse: morphologie du dernier stade larvaire, de la puppe et notes biologiques (Diptera, Syrphidae). [*Mallota cimbiciformis* (Fallén), new for the Swiss fauna: last instar larva, puparium and notes on the biology (Diptera, Syrphidae)]. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 62: 67–78.
- Maibach A. & Goeldlin de Tiefenau P. 1992. Description de la puppe de *Chalcosyrphus (Xylotodes) eunotus* (Loew) et synthèse caractéristiques morphologiques des stades immatures de plusieurs genres de la tribu des Xylotini (Diptera, Syrphidae). [Description of the pupa of *Chalcosyrphus (Xylotodes) eunotus* (Loew) and summary of the morphological characteristics of the immature stages of several genera of the Xylotini tribe (Diptera, Syrphidae)]. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 65: 165–175.
- Marcos-García M^a.A., Micó E., Quinto J., Briones R. & Galante E. 2011. Lo que las oquedades esconden. [What the hollows hide]. *Cuadernos de Biodiversidad* 34: 3–7. [Spanish with English summary]. <https://doi.org/10.14198/cdbio.2011.34.01>
- Martin M., Paillet Y., Larrieu L., Kern C.C., Raymond P., Drapeau P. & Fenton N.J. 2022. Tree-Related Microhabitats are promising yet underused tools for biodiversity and nature conservation: A systematic review for international perspectives. *Frontiers in Forests and Global Change* 5: 818474. <https://doi.org/10.3389/ffgc.2022.818474>
- Martínez-Pérez S., Galante E. & Micó E. 2022. Sex specificity of dispersal behaviour and flight morphology varies among tree hollow beetle species. *Movement Ecology* 10: 41. <https://doi.org/10.1186/s40462-022-00340-7>
- Mei M. 2016. Records of *Tracheliodes varus* (Panzer) and *T. curvitorsus* (Herrich-Schaeffer) from Central Italy (Hymenoptera, Crabronidae). *Ampulex* 8: 16–19. https://www.zobodat.at/pdf/Ampulex_8_0016-0019.pdf
- Miklín J., Hauck D., Konvička O. & Cizek L. 2017. Veteran trees and saproxylic insects in the floodplains of Lower Morava and Dyje rivers, Czech Republic. *Journal of Maps* 13(2): 291–299. <https://doi.org/10.1080/17445647.2017.1300785>
- Miklín J., Sebek P., Hauck D., Konvička O. & Cizek L. 2018. Past levels of canopy closure affect the occurrence of veteran trees and flagship saproxylic beetles. *Diversity and Distributions* 24(2): 208–218. <https://doi.org/10.1111/ddi.12670>
- Moquet L., Laurent E., Bacchetta R. & Jacquemart A-L. 2017. Conservation of hoverflies (Diptera, Syrphidae) requires complementary resources at the landscape and local scales. *Insect Conservation and Diversity* 11(1): 72–87. <https://doi.org/10.1111/icad.12245>
- Morris R. & Ball S. 2021. Death by one hundred droughts: is climate change already driving biodiversity declines in Britain? *British Wildlife* 33: 13–20.
- Myking T., Bøhler F., Austrheim G. & Solberg E.J. 2011. Life history strategies of aspen (*Populus tremula* L.) and browsing effects: a literature review. *Forestry: An International Journal of Forest Research* 84 (1): 61–71. <https://doi.org/10.1093/forestry/cpq044>

- Nicholas S., Thyselius M., Holden M. & Nordström K. 2018. Rearing and long-Term maintenance of *Eristalis tenax* hoverflies for research studies. *Journal of Visualized Experiments* 135: e57711. <https://doi.org/10.3791/57711>
- Nieuwenhuis M. & Barrett F. 2009. The growth potential of downy birch (*Betula pubescens* (Ehrh.)) in Ireland. *Forestry* 75(1): 75–87. <https://doi.org/10.1093/forestry/75.1.75>
- Nunifu T. 2008. Compatible diameter and height increment models for lodgepole pine, trembling aspen, and white spruce. *Canadian Journal of Forest Research* 39(1): 180–192. <https://doi.org/10.1139/X08-168>
- Nol E., Douglas H. & Crins W.J. 2006. Responses of Syrphids, Elaterids and Bees to single-tree selection harvesting in Algonquin provincial park, Ontario. *The Canadian Field-Naturalist* 120(1): 15–21. <https://doi.org/10.22621/cfn.v120i1.239>
- Nolan V., Reader T., Gilbert F. & Atkinson N. 2020. The Ancient Tree Inventory: a summary of the results of a 15 year citizen science project recording ancient, veteran and notable trees across the UK. *Biodiversity and Conservation* 29: 3103–3129. <https://doi.org/10.1007/s10531-020-02033-2>
- Onur Erman D. 2014. Bird Houses in Turkish Culture and Contemporary Applications. *Procedia Social and Behavioral Sciences* 122: 306–311. <https://doi.org/10.1016/j.sbspro.2014.01.1345>
- Orengo-Green J.J., Javier Quinto J., Ricarte A. & Marcos-García, M^a 2023. Combined stereomicroscope and SEM disentangle the fine morphology of the undescribed larva and puparium of the hoverfly *Milesia crabroniformis* (Fabricius, 1775) (Diptera: Syrphidae). *Micron* 165: 103397. <https://doi.org/10.1016/j.micron.2022.103397>
- Pérez-Bañón C., Radenković S., Vujić A., Ståhls G., Rojo S., Grković A. & Petanidou T. 2016. *Brachyopa minima* (Diptera: Syrphidae), a new species from Greece with notes on the biodiversity and conservation of the genus *Brachyopa* Meigen in the Northern Aegean Islands. *Zootaxa* 4072 (2): 217–234. <https://doi.org/10.11646/zootaxa.4072.2.5>
- Peterken G. 2002. Reversing habitat fragmentation British woodlands. WWF-UK, Godalming, Surrey. 60 pp. <https://wwfeu.awsassets.panda.org/downloads/ukforestsfragmentation.pdf>
- Petermann J.S. & Gossner M.M. 2022. Aquatic islands in the sky: 100 years of research on water-filled tree holes. *Ecology and Evolution* 12(8): e9206. <https://doi.org/10.1002/ece3.9206>
- Piovesan G. & Biondi F. 2021. On tree longevity. *New Phytologist* 231(4): 1318–1337. <https://doi.org/10.1111/nph.17148>
- Przepióła F. & Ciach M. 2022. Tree microhabitats in natural temperate riparian forests: An ultra-rich biological complex in a globally vanishing habitat. *Science of the Total Environment* 803: 149881. <https://doi.org/10.1016/j.scitotenv.2021.149881>
- Quinto J., Marcos-García M^a.Á., Díaz-Castelazo C., Rico-Gray V., Brustel H., Galante E. & Micó E. 2012. Breaking down Complex Saproxylic Communities: Understanding Sub-Networks Structure and Implications to Network Robustness. *PLoS ONE* 7(9): e45062. <https://doi.org/10.1371/journal.pone.0045062>
- Ramón Arévalo J., Encina-Domínguez J.A., Mellado M., García-Martínez J.E. & Cruz-Anaya A. 2021. Impact of 25 years of grazing on the forest structure of *Pinus cembroides* in northeast Mexico. *Acta Oecologica* 111: 103743. <https://doi.org/10.1016/j.actao.2021.103743>
- Read H. 2000. Veteran trees and guide to good management. Veteran Tree Initiative. 93 pp. <https://ancienttreeforum.co.uk/wp-content/uploads/2015/03/Veteran-Trees-A-Guide-to-Good-Management-almost-complete.pdf>
- Renema W. 2000. Vliegen vliegen? [Do fly fly?]. *Zweefvliegen Nieuwsbrief* 4 (2): 4–95. <https://natuurtijdschriften.nl/pub/555939/ZVNB2000004002003.pdf>

- Ricarte A., Marcos-García M^a.Á., Pérez-Bañón C. & Rotheray G.E. 2007. The early stages and breeding sites of four rare saproxylic hoverflies (Diptera: Syrphidae) from Spain. *Journal of Natural History* 41 (25–28): 1717–1730. <https://doi.org/10.1080/00222930701495046>
- Ricarte A., Souba-Dols G.J., Hauser M. & Marcos-García M^a.Á. 2017. A review of the early stages and host plants of the genera *Eumerus* and *Merodon* (Diptera: Syrphidae), with new data on four species. *PLoS ONE* 12(12): e0189852. <https://doi.org/10.1371/journal.pone.0189852>
- Rösch M. 2012. Forest, Wood, and Ancient Man. *Interdisciplinaria archaeologica Natural Sciences in Archaeology* 3(2): 247–255. <https://doi.org/10.24916/iansa.2012.2.7>
- Rotheray E.L. 2013. Differences in ecomorphology and microhabitat use of four saproxylic larvae (Diptera, Syrphidae) in Scots pine stump rot holes. *Ecological Entomology* 38(3): 219–229. <https://doi.org/10.1111/een.12009>
- Rotheray E.L., Bussiere L.F., Moore P., Bergström L. & Goulson D. 2014. Mark recapture estimates of dispersal ability and observations on the territorial behaviour of the rare hoverfly *Hammerschmidtia ferruginea* (Diptera, Syrphidae). *Journal of Insect Conservation* 18(2): 179–188. <https://doi.org/10.1007/s10841-014-9627-7>
- Rotheray E.L. & MacGowan I. 2015. Pine hoverfly. Version 1.0. *In*: Gaywood M.J., Boon P.J., Thompson D.B.A. & Strachan I.M. (eds). *The Species Action Framework Handbook*, Scottish Natural Heritage, Battleby, Perth.
- Rotheray E.L., MacGowan I., Rotheray G.E., Sears J. & Elliott A. 2008. The conservation requirements of an endangered hoverfly, *Hammerschmidtia ferruginea* (Diptera, Syrphidae) in the British Isles. *Journal of Insect Conservation*. 13(6): 569–574. <https://doi.org/10.1007/s10841-008-9204-z>
- Rotheray G.E. 1991. Larval stages of 17 rare and poorly known British hoverflies (Diptera: Syrphidae). *Journal of Natural History* 25(4): 945–969. <https://doi.org/10.1080/00222939100770621>
- Rotheray G.E. 1993. A Colour guide to hoverfly larvae in Britain and Europe. *Dipterists Digest* 9: 1–155.
- Rotheray G.E. 2004. Autecology and conservation of *Callicera spinolae* the golden hoverfly (Diptera, Syrphidae). *English Nature Research Reports* 581: 1–40. <http://publications.naturalengland.org.uk/publication/133001>
- Rotheray G.E. & Gilbert F.S. 2011. *The Natural History of Hoverflies*. Forrest Text, Cardigan, UK. 334 pp.
- Rotheray G.E., Hancock E.G., Hewitt S., Horsfield D., MacGowan I., Robertson D. & Watt K. 2001. The biodiversity and conservation of saproxylic Diptera in Scotland. *Journal of Insect Conservation* 5: 77–85. <https://doi.org/10.1023/A:1011329722100>
- Rotheray G.E. & MacGowan I. 2000. Status and breeding sites of three presumed endangered Scottish saproxylic syrphids (Diptera, Syrphidae). *Journal of Insect Conservation* 4: 215–223. <https://doi.org/10.1023/A:1011380316156>
- Rotheray G.E. & Perry I. 1994. The larva of *Callicera spinolae* with a key to the larvae of British *Callicera* species (Diptera, Syrphidae). *The Entomologist* 113 (3&4): 205–210.
- Rotheray G.E. & Rotheray E.L. 2012. Translocating the Pine Hoverfly, *Blera fallax*. *Antenna* 36(1): 36–41.
- Rotheray G.E. & Stuke J-H. 1998. Third stage larvae of four species of saproxylic Syrphidae (Diptera), with a key to the larvae of British *Criorhina* species. *Entomologist's Gazette* 49: 209–217.
- Rubino D. & McCarthy B. 2003. Composition and ecology of macrofungal and myxomycete communities on oak woody debris in a mixed-oak forest of Ohio. *Canadian Journal of Forest Research* 33(11): 2151–2163. <https://doi.org/10.1139/x03-137>

- Ruegger N. 2016. “Bat Boxes - A Review of Their Use and Application, Past, Present and Future.” *Acta Chiropterologica* 18(1): 279–299.
<https://doi.org/10.3161/15081109ACC2016.18.1.017>
- RZSS 2022. Conservation efforts establish new populations of critically endangered insect in Britain. <https://www.rzss.org.uk/news/article/21195/conservation-efforts-establish-new-populations-of-critically-endangered-insect-in-britain--> [accessed on 2–11–2022]
- Sánchez-Galván I.R., Marcos-García M^a.Á., Galante E., Azeria E.T. & Micó E. 2014. Unraveling Saproxylic Insect Interactions in Tree Hollows from Iberian Mediterranean Forest. *Environmental Entomology* 47(2): 300–308. <https://doi.org/10.1093/ee/nvy008>
- Sandström J., Bernes C., Junninen K., Löhmus A., Macdonald E., Müller J. & Jonsson B.G. 2019. Impacts of dead wood manipulation on the biodiversity of temperate and boreal forests. A systematic review. *Journal of Applied Ecology* 56(7): 1770–1781.
<https://doi.org/10.1111/1365-2664.13395>
- Schlaghamerský D.J. 2003. Saproxylic invertebrate of floodplains, a particularly endangered component of biodiversity. *In*: F. Mason F., Nardi G. & Tisato M. (eds). *Dead wood: a key to biodiversity. Proceedings of the International Symposium 29–31 May 2003.* Mantova Italy, 99 pp.
- Schlaghamerský J. & Omelková M. 2007. The present distribution and nest tree characteristics of *Liometopum microcephalum* (Panzer, 1798) (Hymenoptera: Formicidae) in South Moravia. *Myrmecological News* 10: 85–90.
- Sebek P., Altman J., Platek M. & Cizek L. 2013. Is active management the key to the conservation of saproxylic biodiversity? Pollarding promotes the formation of tree hollows. *PLoS ONE* 8(3): e60456. <https://doi.org/10.1371/journal.pone.0060456>
- Sist P., Mazzei L., Blanc L. & Rutishauser E. 2014. Large trees as key elements of carbon storage and dynamics after selective logging in the Eastern Amazon. *Forest Ecology and Management* 318: 103–109. <https://doi.org/10.1016/j.foreco.2014.01.005>
- Sjuts B. 2004. Das Vorkommen von *Brachyopa insensilis* Collin, 1939 (Diptera, Syrphidae) an *Aesculus hippocastanum* L. (Sapindales, Hippocastanaceae) in der Stadt Leer. *Volucella* 1: 211–215. https://www.zobodat.at/pdf/Volucella_7_0211-0215.pdf
- Schmid U. & Moertelmaier T. 2007. The larvae of *Brachypalpus chrysites* Egger, 1859 and *Chalcosyrphus valgus* (Gmelin, 1790) (Diptera, Syrphidae). *Volucella* 8: 109–120.
https://www.zobodat.at/pdf/Volucella_8_0109-0120.pdf
- Speight M.C.D. 1989. Saproxylic invertebrates and their conservation. Strasbourg (France). *Nature and Environment Series* 42: 1–79.
- Soszyńska-Maj A., Soszyński B. & Klasa A. 2009. Distribution and ecology of the saproxylic hoverfly *Chalcosyrphus eunotus* (Loew, 1873) (Diptera: Syrphidae) in Poland. *Fragmenta Faunistica* 52 (2): 191–195. http://rcin.org.pl/Content/43842/PDF/WA058_53445_P256-T52_Frag-Faun-Nr-2.pdf
- Ssymank A. 1991. Die funktionale Bedeutung des Vegetationsmosaiks eines Waldgebietes der Schwarzwaldvorbergzone für blütenbesuchende Insekten - untersucht am Beispiel der Schwebfliegen (Diptera, Syrphidae). *Phytocoenologia (Stuttgart-Lehre)* 19(3): 307–390.
<http://doi.org/10.1127/phyto/19/1991/307>
- Ssymank A. 2001. Vegetation und blütenbesuchende Insekten in der Kulturlandschaft. – Pflanzengesellschaften, Blühphänologie, Biotopbindung und Raumnutzung von Schwebfliegen (Diptera, Syrphidae) im Drachenfelder Ländchen sowie Methodenoptimierung und Landschaftsbewertung. *Tierwelt in der Zivilisationslandschaft, Teil V. Schriftenr. Landschaftspfl. u. Naturschutz* 64, 513 pp. Bonn-Bad Godesberg.
- Ssymank A. & Gilbert F.S. 2008. Anemophilous pollen in the diet of Syrphid flies with special reference to the leaf feeding strategy occurring in Xylotini. (Diptera, Syrphidae).

- Deutsche Entomologische Zeitschrift 40(2): 245–258.
<https://doi.org/10.1002/mmnd.19930400204>
- Ssymank A., Kearns C.A., Pape T. & Thompson F.C. 2008. Pollinating Flies (Diptera): A major contribution to plant diversity and agricultural production. *Biodiversity* 9 (1–2): 86–89. <https://doi.org/10.1080/14888386.2008.9712892>
- Stiven R. 2009. Management of ancient wood pasture. Forestry Commission Scotland, Edinburgh, UK. 14 pp.
<https://forestry.gov.scot/images/corporate/pdf/fcsancientwoodpastureguidance.pdf>
- Tăușan I. 2018. Updated distribution of the velvety tree ant *Liometopum microcephalum* (Panzer, 1798) (Hymenoptera: Formicidae) in Romania. *Brukenthal Acta Musei*, XIII. 3: 449–52.
- Taylor H., Rotheray E.L., Elliot A., MacGowan I., Sears J. & Tompkins G. 2021. Hovering on the edge of extinction: efforts to save the Pine Hoverfly. *British Wildlife* 32(8): 547–554.
- Thorn, S., Seibold S., Leverkus A.B., Michler T., Müller J., Noss R.F., Stork N., Vogel S. & Lindenmayer D.B. 2020. The living dead: acknowledging life after tree death to stop forest degradation. *Frontiers in Ecology and the Environment* 18(9): 505–512.
<https://doi.org/10.1002/fee.2252>
- Ulyshen M.D. 2018. Saproxylic insects: Diversity, ecology and conservation. Springer International Publishing. 904 pp. <https://doi.org/10.1007/978-3-319-75937-1>
- van Eck A. 2016. *Sphegina*-larven (Diptera, Syrphidae) in de bronbossen van Beek-Ubbergen. [*Sphegina* larvae (Diptera, Syrphidae) in the source forests of Beek-Ubbergen]. *De Vliegenmepper* 25(2): 12–16.
- van Eck A., Heitmans W.R.B. & Polaszek A. 2016. *Tetrastichus brachyopae* (Hymenoptera: Eulophidae) new to the Netherlands, reared from *Brachyopa* larvae (Diptera: Syrphidae). *Entomologische Berichten* 76 (6): 226–230.
<https://secties.nev.nl/pages/publicaties/eb/nummers/2016/76-6/226-230.pdf>
- van Rijn P.C.J. & Wäckers F.L. 2016. Nectar accessibility determines fitness, flower choice and abundance of hoverflies that provide natural pest control. *Journal of Applied Ecology* 53: 925–933. <https://doi.org/10.1111/1365-2664.12605>
- van Steenis J. 1989. Zweefvliegenlarven in een boomgaard. *Protter* 56: 16–17.
- van Steenis J. 2015. Syrphidae in trees; artificial breeding sites for saproxylic Syrphidae. Poster presentation 8th International Symposium on Syrphidae 4–8th June 2015 Monschau, Germany.
- van Steenis J. 2016. The hoverfly (Diptera: Syrphidae) fauna of the nature reserve Hågadalen-Nåsten, Uppsala, Sweden. *Entomologisk Tidskrift* 137 (3): 111–129.
http://www.sef.nu/download/entomologisk_tidskrift/et_vol_137_2016/ET-2016-111-129.pdf
- van Steenis J. 2022. Syrphidae in Trees, JvS members.
<https://www.syrphidaeintrees.com/new-syrphidae/> [accessed 23-11-2022]
- van Steenis J., van Steenis W. & Wakkie B. 2001. Hoverflies in southern Skåne, Sweden (Diptera: Syrphidae). *Entomologisk Tidskrift* 122 (1–2): 15–27.
[https://www.sef.nu/download/entomologisk_tidskrift/et_2001/ET2001%2015-27\(2\).pdf](https://www.sef.nu/download/entomologisk_tidskrift/et_2001/ET2001%2015-27(2).pdf)
- van Steenis J., Nedeljković Z., Tot T., Ent L.J., van der Eck A., Mazánek L., Šebić A., Radenković S. & Vujić A. 2019. New records of hoverflies (Diptera: Syrphidae) and the rediscovery of *Primocerioides regale* Violenitsh for the fauna of Serbia. *Biologica Serbica* 41(1): 94–103. <https://doi.org/10.5281/zenodo.3526446>
- van Steenis J., van Zuijen M.P., Bot S., van der Ent L.-J., Barkalov A., van Eck A., Fleury J., Földesi R., Heimburg H., Hadrava J., Koch B., Lutovinovas E., Mazanek L., Van de Meutter F., Mielczarek Ł., Palmer C.J., Popov G.V., Radenković S., Reemer M.,

- Ssymank A.M., van Steenis W., Tóth S., Vujić A. & Wakkie B. 2020. Faunistical overview of the European species of the genera *Brachyopa* Meigen, 1822 and *Hammerschmidtia* Schummel, 1834 (Diptera, Syrphidae). *Bonn zoological Bulletin* 69(2): 309–366. <https://doi.org/10.20363/BZB-2020.69.2.309>
- Vujić A., Gilbert F., Flinn G., Englefield E., Ferreira C.C., Varga Z., Eggert F., Woolcock S., Böhm M., Mergy R., Ssymank A., van Steenis W., Aracil A., Földesi R., Grković A., Mazanek L., Nedeljković Z., Pennards G.W.A., Pérez C., Radenković S., Ricarte A., Rojo S., Ståhls G., van der Ent L.-J., van Steenis J., Barkalov A., Campoy A., Janković M., Likov L., Lillo I., Mengual X., Milić D., Miličić M., Nielsen T., Popov G., Romig T., Šebić A., Speight M., Tot T., van Eck A., Veselić S., Andric A., Bowles P., De Groot M., Marcos-García M.A., Hadrava J., Lair X., Malidžan S., Nève G., Obreht Vidakovic D., Popov S., Smit J.T., Van De Meutter F., Veličković N. & Vrba J. 2022. Pollinators on the edge: our European hoverflies. *The European Red List of Hoverflies*. Brussels, Belgium: European Commission. 108 pp. <https://wikis.ec.europa.eu/download/attachments/23462140/European%20Red%20List%20of%20Hoverflies.pdf?version=1&modificationDate=1665404637685&api=v2>
- Vujić A., Miličić M., Janković Milosavljević M., van Steenis J., Macadam C., Raser J. & Hochkirch A. in press. Hoverflies specialized to veteran trees in Europe: Conservation Action Plan 2023-2030. IUCN, Brussels. 50 pp.
- Wetherbee R., Birkemoe T. & Sverdrup-Thygeson. A. 2020. Veteran trees are a source of natural enemies. *Nature Scientific Reports* 10: 18485 <https://doi.org/10.1038/s41598-020-75723-0>
- Wetherbee R., Birkemoe T., Burner R.C. & Sverdrup-Thygeson A. 2021. Veteran trees have divergent effects on beetle diversity and wood decomposition. *PLoS One* 16(3): e0248756. <https://doi.org/10.1371/journal.pone.0248756>
- Wolton R. & Luff M. 2016. Observations on the Diptera and other insects frequenting sap exudations on an oak tree in Devon, south-west England. *Dipterists Digest* 23(2): 119–136. <https://dipterists.org.uk/sites/default/files/pdf/Dipterists%20Digest%202016%20Vol%2023%20No%202.pdf>